

I U C L I D

Data Set

Existing Chemical : ID: 88-74-4
CASNo. : 88-74-4
EINECS Name : 2-nitroaniline
EINECS No. : 201-855-4
TSCA Name : Benzenamine, 2-nitro-
Molecular Formula : C6H6N2O2

Producer Related Part
Company : Solutia Inc.
Creation date : 04.04.2002

Substance Related Part
Company : Solutia Inc.
Creation date : 04.04.2002

Memo :

Printing date : 07.11.2002
Revision date :
Date of last Update : 07.11.2002

Number of Pages : 43

Chapter (profile) : Chapter: 1, 2, 3, 4, 5, 7
Reliability (profile) : Reliability: without reliability, 1, 2, 3, 4
Flags (profile) : Flags: without flag, confidential, non confidential, WGK (DE), TA-Luft (DE),
Material Safety Dataset, Risk Assessment, Directive 67/548/EEC, SIDS

1.0.1 OECD AND COMPANY INFORMATION

1.0.2 LOCATION OF PRODUCTION SITE

1.0.3 IDENTITY OF RECIPIENTS

1.1 GENERAL SUBSTANCE INFORMATION

1.1.0 DETAILS ON TEMPLATE

1.1.1 SPECTRA

1.2 SYNONYMS

1.3 IMPURITIES

1.4 ADDITIVES

1.5 QUANTITY

1.6.1 LABELLING

1.6.2 CLASSIFICATION

1.7 USE PATTERN

1.7.1 TECHNOLOGY PRODUCTION/USE

1.8 OCCUPATIONAL EXPOSURE LIMIT VALUES

1.9 SOURCE OF EXPOSURE

1.10.1 RECOMMENDATIONS/PRECAUTIONARY MEASURES

1.10.2 EMERGENCY MEASURES

1.11 PACKAGING

1.12 POSSIB. OF RENDERING SUBST. HARMLESS

1.13 STATEMENTS CONCERNING WASTE

1.14.1 WATER POLLUTION

1.14.2 MAJOR ACCIDENT HAZARDS

1.14.3 AIR POLLUTION

1.15 ADDITIONAL REMARKS

1.16 LAST LITERATURE SEARCH

1.17 REVIEWS

1.18 LISTINGS E.G. CHEMICAL INVENTORIES

2.1 MELTING POINT

Value : = 71.5 °C
Sublimation :
Method : other
Year : 1989
GLP : no data
Test substance : no data
Test substance : Technical grade ONA had purity of > 99% and was likely the source used.
Reliability : (2) valid with restrictions
Listed as Peer Reviewed reference in Hazardous Substances Data Bank (2002) for 2-nitroaniline.
Flag : Critical study for SIDS endpoint
24.10.2002 (4)

2.2 BOILING POINT

Value : = 284 °C at
Decomposition :
Method : other
Year : 1989
GLP : no data
Test substance : no data
Reliability : (2) valid with restrictions
Listed as Peer Reviewed reference in Hazardous Substances Data Bank (2002) for 2-nitroaniline.
Flag : Critical study for SIDS endpoint
24.10.2002 (4)

2.3 DENSITY

2.3.1 GRANULOMETRY

2.4 VAPOUR PRESSURE

Value : = .0368 hPa at 25° C
Decomposition :
Method : other (calculated)
Year : 1989
GLP : no data
Test substance : no data
Reliability : (2) valid with restrictions
Cited as Peer Reviewed reference in Hazardous Substances Data Bank (2002) for 2-nitroaniline.
Flag : Critical study for SIDS endpoint
24.10.2002 (5)

2.5 PARTITION COEFFICIENT

Log pow : 1.85 at °C
Method : other (calculated)

2. Physico-Chemical Data

Id 88-74-4
Date 07.11.2002

Year : 1985
GLP : no data
Test substance : no data
Reliability : (2) valid with restrictions
Listed as Peer Reviewed reference in Hazardous Substances Data Bank (2002) for 2-nitroaniline and listed as Recommended value in SRC CHEMFATE data base (2002).
Flag : Critical study for SIDS endpoint
24.10.2002 (8)

2.6.1 WATER SOLUBILITY

Value : = 1470 mg/l at 25 ° C
Qualitative :
Pka : at 25 ° C
PH : at and ° C
Method : other
Year : 1991
GLP : no data
Test substance : no data
Reliability : (2) valid with restrictions
Listed as Peer Reviewed reference in Hazardous Substances Data Bank (2002) for 2-nitroaniline and SRC CHEMFATE Data base (2002).
Flag : Critical study for SIDS endpoint
24.10.2002 (18)

2.6.2 SURFACE TENSION

2.7 FLASH POINT

2.8 AUTO FLAMMABILITY

2.9 FLAMMABILITY

2.10 EXPLOSIVE PROPERTIES

2.11 OXIDIZING PROPERTIES

2.12 ADDITIONAL REMARKS

3.1.1 PHOTODEGRADATION

Type	: air
Light source	: other
Light spect.	: > 290 nm
Rel. intensity	: based on Intensity of Sunlight
Direct photolysis	
Half-life t _{1/2}	: = 9.5 hour(s)
Degradation	: = 16 % after 3 hour(s)
Quantum yield	:
Indirect photolysis	
Sensitizer	: OH
Conc. of sens.	:
Rate constant	: = .000000000013 cm ³ /(molecule*sec)
Degradation	: % after
Deg. Product	:
Method	: other (calculated)
Year	: 2002
GLP	: no
Test substance	: no data
Method	: Direct photodegradation measured using a medium-pressure mercury arc emitting > 290 mu; irradiations were conducted in triethylamine for 3 hrs; Additionally, a calculated value of 9.5 hr was derived by AOP Computer program v1.90. The program estimates the Atmospheric Oxidation Potential by estimating the rate constant for the atmosphere, gas-phase reaction between photochemically produced hydroxyl radicals and organic chemicals. The methodology is based on the SAR methods developed by Atkinson et al, 1987, Intern. J. Chem. Kinet. 19: 799-828 and described by Meylan and Howard, 1993, Chemosphere 26:2293-2299.
Reliability	: (2) valid with restrictions Measurements published in a peer reviewed journal. Estimated value based on model recommended by US EPA.
Flag	: Critical study for SIDS endpoint
25.10.2002	(1)

3.1.2 STABILITY IN WATER

3.1.3 STABILITY IN SOIL

3.2 MONITORING DATA

3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

Type	: fugacity model level III
Media	: other
Air (level I)	: .525
Water (level I)	: 36.1
Soil (level I)	: 63.2
Biota (level II / III)	:
Soil (level II / III)	: .111
Method	: other
Year	: 2002
Method	: Estimation using measured values from reference documents were possible and incorporated into EPIWIN from Syracuse Research Corp and

3. Environmental Fate and Pathways

Id 88-74-4

Date 07.11.2002

Results

possible and incorporated into EPIWIN from Syracuse Research Corp and based on Meylan, 1993 methodology as adopted by Mackay et al 1996. Second Soil entry includes data in Sediments. Values employed were: Mol wt=138.13; Gebrt;s KC=5.9e-008 atm-m3/mole (Henry database); Vapor Press=0.00277 mm Hg (user entry); Log Kow=1.85 (user entry); Soil Koc=29 (calc by model). Emissions rates for each of the three compartments (air, soil and water) were 1000 kg/hr.

Chem Name : o-Nitroaniline

Molecular Wt: 138.13

Henry's LC : 5.9e-008 atm-m3/mole (Henry database)

Vapor Press : 0.00277 mm Hg (user-entered)

Log Kow : 1.85 (user-entered)

Soil Koc : 29 (calc by model)

	Concentration (percent)	Half-Life (hr)	Emissions (kg/hr)		
Air	0.525	19.1	1000		
Water	36.1	900	1000		
Soil	63.2	900	1000		
Sediment	0.111	3.6e+003	0		

	Fugacity (atm)	Reaction (kg/hr)	Advection (kg/hr)	Reaction (percent)
Advection (percent)				
Air	2.03e-011	418	115	13.9
Water	1.69e-012	609	791	20.3
Soil	3.3e-011	1.07e+003	0	35.6
Sediment	1.53e-012	0.469	0.0487	0.0156

Persistence Time: 730 hr

Reaction Time: 1.05e+003 hr

Advection Time: 2.42e+003 hr

Percent Reacted: 69.8

Percent Advected: 30.2

Half-Lives (hr), (based upon Biowin (Ultimate), several screening studies

showing poor biodegradation and Aopwin):

Air: 19.08

Water: 900

Soil: 900

Sediment: 3600

Biowin estimate: 2.589 (weeks-months)

Advection Times (hr):

Air: 100

Water: 1000

Sediment: 5e+004

Reliability

: (2) valid with restrictions

Estimated values based on model recommended by US EPA.

Flag

24.10.2002

: Critical study for SIDS endpoint

(6)

3.3.2 DISTRIBUTION

3.4 MODE OF DEGRADATION IN ACTUAL USE

3. Environmental Fate and Pathways

Id 88-74-4
Date 07.11.2002

3.5 BIODEGRADATION

Type : aerobic
Inoculum :
Concentration : 5mg/l related to Test substance
related to
Contact time : 24 hour(s)
Degradation : = 7 % after 10 month
Result : under test conditions no biodegradation observed
Deg. Product :
Method : other
Year : 1975
GLP : no
Test substance : other TS
Method : Semi-continuous activated sludge (SCAS) test was carried out over a 10-month period at a final addition rate of 5 mg ONA per 24-hr cycle. The methodology used was a standard procedure published in JAOCS 42:986 (1965) and used the modified feed techniques as described in JAOCS 46:432 (1969). ONA concentration was determined using UV spectrophotometry after extraction of the sludge with methylene chloride. Analysis was performed on one 24-hr cycle per week. Activated sludge obtained from local waste treatment facility.
Result : No significant biodegradation occurred, as a mean (+/-95% CI) loss was 7 (+/-11) %. No evidence of any inhibition of the normal sludge growth rate was observed.
Test substance : Used Technical grade ONA with purity > 99%.
Reliability : (2) valid with restrictions
Study was conducted prior to codification of GLPs but is considered well documented. The methodology used in this study has now been codified into internationally accepted test guidance for biodegradability determination.
Flag : Critical study for SIDS endpoint
24.10.2002 (16)

3.6 BOD5, COD OR BOD5/COD RATIO

3.7 BIOACCUMULATION

3.8 ADDITIONAL REMARKS

4.1 ACUTE/PROLONGED TOXICITY TO FISH

Type	: semistatic
Species	: Brachydanio rerio (Fish, fresh water)
Exposure period	: 96 hour(s)
Unit	: mg/l
Analytical monitoring	: yes
LC50	: = 19.5
Method	: Directive 84/449/EEC, C.1 "Acute toxicity for fish"
Year	: 1991
GLP	: no data
Test substance	: other TS
Method	: 96 hr acute toxicity test was conducted in a semistatic system according to the OECD Guideline 202, as published in 1984. Zebrafish were approx. 3 mo. of age and weighed between 200-350 mg; both sexes were used. Fish were not fed 24h prior to testing and during the 96-h exposure period. A 12-h light;dark cycle was employed. The test water was charcoal-filtered, aerated tap water which was mixed with a stock solution of the chemical in distilled water and stirred at room temperature. The pH, dissolved oxygen and temperature of the water were 8.6+/-0.3, 85+/_15% and 26.5+/-1 degree C., respectively. Once a day the concentrations were checked photometrically and the test solutions were renewed if required. LC50 values were calculated using a computer program based on the method of Litchfield and Wilcoxon (1949).
Result	: The 96 hr LC50 was determined to be 19.5 mg/l with SE of +/- 1.7 mg/L.
Test substance	: Test sample purchased from a chemical supplier; Technical grade was typically > 99%.
Reliability	: (1) valid without restriction No information was reported in the article about conduct under GLPs; however, as this study was conducted specifically to meet OECD test guideline 202 it is reasonable to assume that it was conducted under GLPs as well.
Flag	: Critical study for SIDS endpoint
16.10.2002	(19)

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Type	: static
Species	: Daphnia magna (Crustacea)
Exposure period	: 48 hour(s)
Unit	: mg/l
Analytical monitoring	:
NOEC	: >= 12.5
EC50	: = 14.5
Method	: EPA OTS 797.1300
Year	: 1983
GLP	: yes
Test substance	: other TS
Method	: Test article dissolved in Dimethyl Formamide (0.5 ml/L) and introduced to glass jars filled with well water; DO, pH, alkalinity and hardness measured prior to and after testing. Three replicates run, using 10 Daphnia per dosage level per rep. Dosages evaluated: control, solvent control, 6.25, 12.5, 25, 50 and 100 mg/L.
Result	: EC50 values (95% CI) of 18.7 (12.5-25) mg/L at 24 hr and 14.5 (12.5-25) mg/L. at 48-hr interval. The NOEC was 12.5 mg/L. Following was the % deaths observed: At 24 hr- Control (0%), solvent control (0 %), 6.25 mg/L (0 %), 12.5 (0 %), 25 (0 %), 50 (93.3%) and 100 mg/L (100%); At 48 hr -

	Control (0%), solvent control (0%), 6.25 mg/L (0 %), 12.5 (30%), 25, (100%), 50 (100%), and 100 mg/L (100%). pH and dissolved oxygen ranged from 7.0-8.4 and 7.8-9.3 mg/L, respectively. The mean temp. was 23.7 degrees C. Alkalinity ranged between 298-400 mg/L and water hardness ranged between 220-370 mg/L. Evidence of insolubility of test substance was seen at 100 mg/L.	
Test substance	: Used Technical grade ONA, with purity of > 99%.	
Reliability	: (1) valid without restriction Study conducted according to ASTM/EPA guidance, which is consistent with OECD test guidance.	
Flag	: Critical study for SIDS endpoint	
16.10.2002		(15)

4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

Species	: Scenedesmus sp. (Algae)	
Endpoint	: growth rate	
Exposure period	: 48 hour(s)	
Unit	: mg/l	
Analytical monitoring	: no data	
EC50	: = 64.5	
Method	: OECD Guide-line 201 "Algae, Growth Inhibition Test"	
Year	: 2001	
GLP	: no data	
Test substance	: other TS	
Method	: A 48-hr algae inhibition test following OECD test methods was conducted using <i>S. obliquus</i> as the test organism. Five concentration gradients were used, in concentration spacing of 0.2. pH of the culture medium was adjusted to 7.2+/-0.2. Two replicates of each concentration and untreated control were run. The algae in the logarithmic growing period were inoculated into 250 ml Erlenmeyer flasks, and added to 60 ml of the culture media, compound and algae. The initial algae cell concentration was approx. 1 x 10E4 cells/ml. The culture was incubated under a continuous light by fluorescent bulb at 20+/-1 degree C and average illumination intensity of 4000 lux. Growth was monitored by electron microscope (400X). EC values were determined by one variable linear regression analysis.	
Test substance	: Test sample purchased from chemical supplier; typical technical grade purity of ONA was 99%.	
Reliability	: (1) valid without restriction No mention made regarding conduct under GLPs in article; however, as this study was conducted specifically to meet OECD guideline 201 it can reasonably be assumed that it also was conducted under GLPs.	
Flag	: Critical study for SIDS endpoint	
27.08.2002		(7)

4.4 TOXICITY TO MICROORGANISMS E.G. BACTERIA

4.5.1 CHRONIC TOXICITY TO FISH

4.5.2 CHRONIC TOXICITY TO AQUATIC INVERTEBRATES

4.6.1 TOXICITY TO SOIL DWELLING ORGANISMS

4.6.2 TOXICITY TO TERRESTRIAL PLANTS

4.6.3 TOXICITY TO OTHER NON-MAMM. TERRESTRIAL SPECIES

4.7 BIOLOGICAL EFFECTS MONITORING

4.8 BIOTRANSFORMATION AND KINETICS

4.9 ADDITIONAL REMARKS

5.1.1 ACUTE ORAL TOXICITY

Type : LD50
Species : rat
Strain : Sprague-Dawley
Sex : male/female
Number of animals : 20
Vehicle : other
Value : = 2050 mg/kg bw
Method : other
Year : 1977
GLP : no
Test substance : other TS
Method : calc. method of deBeer, 1945, J. Pharmacol. Experimen. Ther. 85:1.
 Test substance was Technical grade ONA with purity of > 99%; administered as 10% corn oil solution
 Used 5 rats (mixed sex) /group. Four groups of rats were administered test article by gavage in increasing doses at increments of 0.1 fractional log intervals. Clinical signs recorded daily and body wts. recorded weekly. Animals observed for 14 days. Necropsies were performed on all animals. Food and water given ad libitum; humidity and temp. controlled.

Result : OLD50=2050 mg/kg; 95% CI of 1760-2380; all deaths occurred within 24 hrs.; Deaths: 1260-0/5; 1580-1/5, 2000-2/5, 2510-5/5; Signs of toxicity: yellow colored urine, generalized weakness; Observations at autopsy for decedents-hemorrhagic lungs, liver hyperemia, abdominal cavity yellow stained, g.i. irritation; for survivors - viscera appeared normal.

Reliability : (2) valid with restrictions
 Conducted using fewer animals than # 401; conduct consistent with but prior to enactment of GLP guidelines; This was a supplemental study to the HPV program in that an acute study by another route has been used to fulfill this HPV data endpoint.

07.11.2002

(17)

5.1.2 ACUTE INHALATION TOXICITY

Type : LC0
Species : rat
Strain : Wistar
Sex : male/female
Number of animals : 10
Vehicle : other
Exposure time : 4 hour(s)
Value : > 2529 mg/m³
Method : OECD Guide-line 403 "Acute Inhalation Toxicity"
Year : 1996
GLP : yes
Test substance : other TS
Method : Test article used was 65% aqueous solution of Technical grade ONA (typical purity of 99%). Groups of 5 male and 5 female rats were exposed to a single aerosol concentration of ONA solution in PEG (to facilitate nebulization) under nose only conditions; the chamber was operated under dynamic exposure conditions. Animals were observed daily for clinical signs; body wts recorded on days 3, 7 and 14. Clinical observations were consistent with a Functional Observational Battery set of indices; methemoglobin determinations were made following exposure. All rats underwent a gross necropsy at study term. Food and water were given ad libitum. Observation period was 14 days. A vehicle control group of rats

was exposed similarly to polyethylene glycol/acetone. Analytical test levels determined by GC method; particle size determined using cascade impactor. Statistical evaluations performed on body weights and physiological data using ANOVA procedures.

Result : Limit test
 No deaths occurred at the maximum achievable level tested of 2,529 mg/m³ (analytical level); the MMAD was 2.1 µm indicating particle sizes of a respirable range. Animals exposed at this level exhibited decrements in body weight gain, hypothermia, distinct discoloration of the urine, and bradypnea, all of which were attributed to test article. These observations persisted no longer than 1 day following exposure. No adverse effects were noted in reflex measurements. No macroscopic findings attributable to test article were observed.

Reliability Flag : (1) valid without restriction
 : Critical study for SIDS endpoint

26.08.2002 (2)

5.1.3 ACUTE DERMAL TOXICITY

Type : LD0
Species : rabbit
Strain : New Zealand white
Sex : male/female
Number of animals : 3
Vehicle : other
Value : > 7940 mg/kg bw
Method : other
Year : 1977
GLP : no
Test substance : other TS
Method : Determination of Minimum Lethal Dose, thus used 1-2 animals /group; 24-hr occlusive dermal patch with 14-day observation period; necropsy at sacrifice, daily cage-side observations made for 2 weeks and weights recorded initially and after 7 and 14 days.
 Test article used was Technical grade ONA with purity > 99%; Administered as 40% solution-suspension in corn oil. Administered to clipped, intact skin of rabbits for 24-hr exposure under occluded conditions. Then removed and animals observed for 14 days.

Result : No deaths (0/1) at 5010 mg/kg or (0/2) at 7940 mg/kg; Observations: Yellow staining, reduced appetite and activity during first 3 days; all normal on day 14. No macroscopic necropsy findings.

Conclusion : Considered sufficient to establish toxicity to rodents by dermal route
Reliability : (2) valid with restrictions
 Used a small no. animals; conducted consistent with but prior to enactment of US GLPs in 1979; this study was a Supplemental study to the HPV program since another study by a another route was chosed to fulfill this HPV Endpoint.

07.11.2002 (17)

5.1.4 ACUTE TOXICITY, OTHER ROUTES

5.2.1 SKIN IRRITATION

5.2.2 EYE IRRITATION

5.3 SENSITIZATION

5.4 REPEATED DOSE TOXICITY

Species	:	rat
Sex	:	male
Strain	:	Sprague-Dawley
Route of admin.	:	inhalation
Exposure period	:	6 hr/day
Frequency of treatment	:	5 days/week for 4 weeks
Post obs. period	:	none
Doses	:	9.8 and 93 mg/m ³ (analytically determined conc.)
Control group	:	yes, concurrent no treatment
NOAEL	:	= 9.8 mg/m ³
LOAEL	:	= 93 mg/m ³
Method	:	OECD Guide-line 412 "Repeated Dose Inhalation Toxicity: 28-day or 14-day Study"
Year	:	1983
GLP	:	yes
Test substance	:	other TS
Method	:	Test material used was Technical grade ONA with purity > 99%. Test article generation used preheated nitrogen which was passed over the test agent in a paraffin oil bath; thus, no solvent, like CELLOSOLV, as used in a previous 4-wk inhalation study (BD-81-322), was employed in this study. This study was designed to determine whether ONA alone induced testicular effects observed in study BD-81-322, using CELLOSOLV solvent; Thus, each test group consisted of 10 male rats; daily observations, hematology (HGB, HCT, RBC, MET, retic, clot time, RBC morph and t/diff. leukocytes) evaluated on all animals prior to sacrifice; Brain and testicular wts were recorded while testes and epididymides were examined grossly and microscopically for all test animals. Body weight, hematology data and absolute and relative organ weights were treated for statistical differences. Parametric analysis was performed using ANOVA methods followed by Dunnet's test when mean differences were observed between dose groups; Kruskal Wallis test and Dunn's rank sum test were used for nonparametric analysis. Both 5% and 1% levels of significance were reported for each parameter. Whole body exposure in stainless steel chamber; analytically determined doses were 9.8 and 93 mg/m ³ respectively. Analysis done by UV 4x daily, particle size confirmed during week 1 and rechecked periodically using Cascade impactor.
Remark	:	This study confirms that ONA produces no effects on testes following inhalation exposure and that results of a previous study (BD-81-322) were the result of use of CELLOSOLV as vehicle. These results, in conjunction with findings in the previous study cited earlier, are sufficient to meet all toxicity parameters established in OECD test guideline 412.
Result	:	Mean testicular wts (absolute and relative) were comparable to controls in both ONA test groups; no gross or microscopic changes in testes/epididymides were observed; Minimal changes in some hematological parameters (increases in methemoglobin i.e. MET and HCT and decreased total leuk. and seg. neutrophils) were seen at 93 mg/m ³
Reliability	:	(1) valid without restriction
Flag	:	Critical study for SIDS endpoint
07.11.2002		
Species	:	rat
Sex	:	male/female
Strain	:	Sprague-Dawley

(11)

5. Toxicity

Id 88-74-4

Date 07.11.2002

Route of admin.	:	inhalation
Exposure period	:	6 hrs/day
Frequency of treatment	:	5 days/week for 4 weeks
Post obs. period	:	none
Doses	:	10, 30 and 73 mg/m ³
Control group	:	yes, concurrent vehicle
NOAEL	:	= 30 mg/m ³
LOAEL	:	= 73 mg/m ³
Method	:	OECD Guide-line 412 "Repeated Dose Inhalation Toxicity: 28-day or 14-day Study"
Year	:	1982
GLP	:	yes
Test substance	:	other TS
Method	:	Test substance used was Technical grade ONA with purity of > 99% which was mixed with 2000 mg/m ³ CELLOSOLVE (ethylene glycol monoethyl ether) as a concurrent vehicle; 10 rats/sex/group were exposed in 1 cub. meter steel/glass chambers via whole body exposure; Analytically determined (4X/d) concentration means were: 10, 27.5 and 73 mg/m ³ , respectively. Particle size means were all below 1 micron for each aerosol concentration. All animals were observed daily for toxic signs, weighed weekly, and underwent examination for clinical chemistries, hematology, ocular toxicity. Organ weights were taken at necropsy and microscopic exams were conducted on over 40 tissues for all high dose and control animals and target organs for all animals. Body weights, food consumption, hematology and clinical chemistry, absolute and relative organ weights were analyzed using ANOVA methods followed by Dunnett's test for parametric parameters while nonparametric parameters were subjected to Kruskal Wallis test followed by Dunn's rank sum test to determine statistical differences. Both 5% and 1% levels of significance were reported for each parameter.
Remark	:	Ambiguous information on testicular effects were resolved with a follow up study (BD-82-270) which assessed the issue of testes effects and the confounding use of Cellosolv as the solvent in this study. Subsequent results confirmed cellosolv as the affective agent.
Result	:	Treatment-related effects : 73 mg/m ³ - Statistically decreased leukocytes in males, and significantly reduced hbg and rbc in females, increased polychromia, anisocytosis and poikilocytosis in males and females, increased rel. liver wts for females (no corresponding histo), decreased absolute and relative testes wts corresponding with degeneration of the germinal epithelium seen microscopically.
Conclusion	:	Study results involving effects on the testes are considered unreliable due to incorrect choice of vehicle control (CELLOSOLVE, which was determined to be a testicular toxin but only after this study was conducted). The issue was resolved after conduct of a follow up study (BD-82-270). However, results in this study confirm that ONA, even in combination with CELLOSOLVE, did not affect measured clinical chemistry parameters, ophthalmology, organ weights, and gross and histopathology of a full set of tissues and organs which were not measured again in the second study (BD-82-270). For this reason, those portions of this study which were indicative of no discernable effect of ONA treatment, can be considered reliable.
Reliability Flag	:	(2) valid with restrictions
16.10.2002	:	Critical study for SIDS endpoint
Species	:	rat
Sex	:	male/female
Strain	:	Sprague-Dawley
Route of admin.	:	gavage
Exposure period	:	14 days

(10)

5. Toxicity

Id 88-74-4

Date 07.11.2002

Frequency of treatment : daily gavage administration throughout test period
Post obs. period : none
Doses : 0, 1, 19, or 100 mg/kg bw
Control group : yes, concurrent vehicle
NOAEL : ≥ 100 mg/kg bw
Method : other
Year : 1989
GLP : no data
Test substance : no data
Method : Groups of 10 M/10 F rats administered test article in corn oil via gavage for 14 consecutive days. A comprehensive evaluation of biochemical, hematological and histopathological evaluations were made at study termination. All animals examined daily for clinical signs and body weights were recorded daily. All animals necropsied on d15 and weights recorded for the following organs: brain, heart, liver, kidney and spleen. Histopathological exams were conducted on approx. 30 tissues and organs, including the gonads. ANOV analyses and Duncan's Multiple Range test ($p < 0.05$) used to determine group differences.

Result : No treatment related findings in hematology, clinical chemistries, clinical observations, body and organ weights or macro- or microscopic findings attributable to treatment

Reliability : (2) valid with restrictions
This study was of insufficient duration to be used to meet HPV testing guidance. It study was provided as Supplemental information as the HPV requirement has been fulfilled with another Repeat Dose study.

07.11.2002

(9)

5.5 GENETIC TOXICITY 'IN VITRO'

Type : Ames test
System of testing : S. typhimurium strains TA98, TA100, TA1535 and TA1537 w & w/o S9
Concentration : 1.5, 3, 6, 7, 15, 30, 40, 150, 225, 450, 600, and 1500 ug/plate
Cycotoxic conc. : 3000 ug/plate (no background lawn) using TA100; 1000 ug/plate tolerated w & w/o S9
Metabolic activation : with and without
Result : negative
Method : Other
Year : 1978
GLP : no
Test substance : other TS
Method : Statistical test used: after data transformation - 1-sided t-test; $p < 0.01$
Test material used was Technical grade ONA with purity of $> 99\%$;
Appropriate positive controls were employed to validate this test methodology.

Result : Negative response seen in spot test at maximum conc. of 10000 ug/plate with and without S9
No significant mutagenic activity seen in any of the 4 tester strains; all positive controls validated adequacy of method used.

Reliability : (2) valid with restrictions
Study conducted consistent with but prior to development of US GLP's in 6/79 and OECD Test Guide 471; study results are confirmed in numerous other published articles.

Flag : Critical study for SIDS endpoint

07.11.2002

(12)

Type : Chromosomal aberration test
System of testing : CHO cells maintained in Eagle MEM media
Concentration : 1 - 10 mM
Cycotoxic conc. : no information provided

Metabolic activation	:	with and without
Result	:	ambiguous
Method	:	other
Year	:	1994
GLP	:	no data
Test substance	:	other TS
Method	:	After overnight incubation in complete medium, the medium was replaced with either serum-free complete medium or an exogenous metabolic activation medium, each containing test material. Cells were treated for 1 h, followed by washing (3X) and incubated in complete medium for either 10h or 16 hr. Colcemid was added for the final 2h of incubation. 100 metaphase cells scored from each of 2 cultures for each treatment level. Negative control group was used. Positive controls included MMS and CP. Statistical package used was EPA's Chromosomal aberration assay data management and analysis system.
Remark	:	This study is Supplemental information as a fully acceptable micronucleus test has been used to fulfill this HPV endpoint.
Result	:	Test material induced a significant ($p < 0.01$) increase in chromosomal aberrations measured 10h after pretreatment both in the presence and absence (1 of 2 trials) of S9. A statistically significant increase in aberrations was also detected after 16h, but only with S9. A dose-response trend was evident in all cases, but only strong responses were observed at the very highest (10 mM) dose tested. The primary aberration observed was a large isochromatid discontinuity seen only in the long arm of the X chromosome. Image enhancement revealed presence of material in the affected region and the alignment of the dislocated segment, making classification of this lesion uncertain. In a separate experiment, all X-chromosome isochromatid anomalies were screened to perform the analysis with and without discontinuity. When excluded, there was no increase in aberrations observed. The cause of this isochromatid discontinuity is uncertain.
Conclusion	:	The authors state that "It is not clear whether this phenomenon represents a legitimate chromosomal aberration."
Reliability	:	(3) invalid
07.11.2002		(3)

5.6 GENETIC TOXICITY 'IN VIVO'

Type	:	Micronucleus assay
Species	:	mouse
Sex	:	male/female
Strain	:	CD-1
Route of admin.	:	i.p.
Exposure period	:	Single doses given twice, 24 hrs apart
Doses	:	0, 50, 250, and 500 mg/kg
Result	:	negative
Method	:	OECD Guide-line 474 "Genetic Toxicology: Micronucleus Test"
Year	:	1989
GLP	:	yes
Test substance	:	other TS
Method	:	Dosages administered in corn oil (10 ml/kg). In a preliminary study, the IP LD50 in mice was determined to be 723 mg/kg; further, the PCD/total erythrocyte ratio was evaluated to determine bone marrow cytotoxicity potential. After completion of dosing, bone marrow was taken from both femors and pooled for slide preparation. Slides were stained with Wright-Giemsa stain pak and scoring was conducted by 2 independent readers. The no. of micronuclear polychromatic erythrocytes (PCEs) per 1000 PCEs and the no. of PCEs and normochromatic erythrocytes/1000 erythrocytes were evaluated for each animal. The individual animal was used as the statistical unit and the Student's T (1-sided) test used to compare treatment

5. Toxicity

Id 88-74-4

Date 07.11.2002

and control group means. A level of $p < 0.05$ was used for all parameters to determine statistical significance.
Highest dosage used was approx. 70% of calc. IP LD50 of 730 mg/kg, as determined in intralaboratory range-find study with mice
Technical grade ONA with purity of > 99% used in this test. Cyclophosphamide (40 mg/kg) positive control used.

Result	:	No increases in micronuclei observed at any ONA dose level; positive control verified the method. Signs of listlessness and unresponsive behavior seen in both sexes at 500 and 250 mg/kg and females at 50 mg/kg ONA; statistically lower body weights observed in females at 500 mg/kg after 48 hr dosing.	
Reliability Flag	:	(1) valid without restriction	
27.08.2002	:	Critical study for SIDS endpoint	(14)
Type	:	Micronucleus assay	
Species	:	mouse	
Sex	:	male/female	
Strain	:	C57BL	
Route of admin.	:	i.p.	
Exposure period	:	Treated twice with 24 h between each treatment	
Doses	:	0, 246, 492 and 738 mg/kg	
Result	:	ambiguous	
Method	:	OECD Guide-line 474 "Genetic Toxicology: Micronucleus Test"	
Year	:	1994	
GLP	:	no data	
Test substance	:	no data	
Method	:	Test article administered IP in olive oil to groups of 5M and 5F mice; controls received only olive oil. High dose reportedly was estimated to be 75% of LD50 as determined in a preliminary experiment. After 36 h following the second treatment, mice were sacrificed and bone marrow removed, a cell suspension made and slides prepared. 500 polychromatic erythrocytes from each animal were scored for the presence of micronuclei. The ratio of PEs to normochromatic cells was also determined to assess cytotoxicity. Data were analyzed using EPA's micronucleus assay data management and analysis system ($p < 0.05$)	
Result	:	No statistically significant increase in PE ratios; thus, no indication of cytotoxicity. A small 1.2 ± 0.08 vs. 2.8 ± 1.50 , but statistically ($p < 0.05$) significant increase in micronuclei was observed at the highest dose tested of 738 mg/kg only in male mice. This effect was observed only in males, not females at this dose level; no effects were seen in either males or females at lower dose levels.	
Reliability	:	(3) invalid	
07.11.2002	:	Considered ambiguous, as the effect noted was small, seen only at one dose level and observed in only one sex. Provided as Supplemental information.	(3)

5.7 CARCINOGENITY

5.8 TOXICITY TO REPRODUCTION

5.9 DEVELOPMENTAL TOXICITY/TERATOGENICITY

Species	:	rat
Sex	:	female
Strain	:	Sprague-Dawley

Route of admin.	: gavage
Exposure period	: Days 6-15 of gestation
Frequency of treatment	: Daily throughout exposure period
Duration of test	: Treated on gestation days 6-15, sacrificed on gestation day 21 for fetal exams
Doses	: 0, 100, 300, 600 mg/kg/day in corn oil
Control group	: yes, concurrent vehicle
NOAEL Maternalt.	: = 100 mg/kg bw
NOAEL Teratogen	: = 600 mg/kg bw
NOAEL Embryotoxicity	: = 600 mg/kg bw
NOAEL Fetotoxicity	: = 600 - mg/kg bw
Method	: OECD Guide-line 414 "Teratogenicity"
Year	: 1985
GLP	: yes
Test substance	: other TS
Method	: 25 pregnant females/group; daily gavage in corn oil at constant volume of 10 ml/kg/d from gestation days 6-15. Dosing solutions were analyzed (GC) for test material concentration and stability periodically throughout the study. Nidation data collected at sacrifice, live fetuses examined externally and by Wilson sections and skeletal exam techniques were used to detect any variations or abnormalities. Body weights and food consumption were collected on gestation days 0, 6, 10, 13, 16 and 21 (day of termination). Daily clinical signs of toxicity recorded on gestation days 6-21. Statistical methods used: body wts. analyzed using Dunnett's test; Counted data (corpora lutea, implants, resorption, live/dead pups) analyzed using Mann-whitney U test; response data (eg. pregnancy rates, litters with postimplantation loss, etc.) assessed with Fischer's exact test and Chi square test.
Result	: Maternal toxicity evidenced by reduced body wt gain at 600 mg/kg and lower food consumption at 600 and 300 mg/kg; both indices were slightly (not stat. signif.) lower than controls, but not considered related to treatment as these events were observed in this group prior to treatment. No effects on pregnancy rates, mean no. live and dead pups, resorptions, nidations, c. lutea; Mean fetal wts were slightly, but not statistically lower than control in 600 mg/kg group. No differences seen in no. litters, fetuses or malformations. One malformation (situs inversus syndrome) was seen in single fetuses from two litters at the 600 mg/kg level; this incidence and lack of correlation to similar findings associated with other mononitroanilines supports the conclusion that this was a spurious finding.
Test substance	: Technical grade of ONA used with purity of > 99%.
Reliability	: (1) valid without restriction
Flag	: Critical study for SIDS endpoint

16.10.2002

(13)

04.04.2002

5.10 OTHER RELEVANT INFORMATION**5.11 EXPERIENCE WITH HUMAN EXPOSURE**

- (1) Bartrop, JA and Bunce, NJ. 1968. Organic photochemistry. Part VIII. The photochemical reduction of nitro-compounds. J. Chem. Soc. (C) 1968:1467-1474.
- (2) Bayer Corp. 1996. Study on acute inhalation toxicity in rats according to OECD No. 403 by T. Martins. Bayer Study no. T3044113. [EPA Document No. 86960000565; Fiche no. OTS0558766]
- (3) Blakey, DH, Maus, KL, Bell, R, Bayley, J, Douglas, GR, and Nestmann, ER. 1994. Mutation Research 320:273-283.
- (4) Budavari, S. (ed.) 1989. The Merck Index- an encyclopedia of chemicals, drugs and biologicals. Whitehouse Station, NJ. p. 1042.
- (5) Daubert, TE and Danner, RP, 1989. Physical and thermodynamic properties of pure chemicals data compilation. Taylor and Francis, Washington, DC.
- (6) EPIWIN, Version, 2002. version 3.10. Syracuse Research Corp, Syracuse, NY.
- (7) G.-H. Lu, Yuan, X, and Zhao, Y.H. 2001. Chemosphere 44:437-440
- (8) Hansch, C and Leo, L, 1985. Medchem Project. Claremont, CA. Pomona College Issue 26.
- (9) Komsta, E, Secours, VE, Chu, I, Valli, VE, Morris, R, Harrison, J, Baranowski, E and Villeneuve, DC. 1989. Bull. Environ. Contam. Toxicol. 43:87-94.
- (10) Solutia study no. BD-81-322. Four week inhalation toxicity study of O-Nitroaniline in the rat. [EPA 878214205; Fiche no. OTS0206486]
- (11) Solutia study no. BD-82-270. Four week inhalation study of Ortho-Nitroaniline in male rats [EPA Document no. 878214205; Fiche no. OTS0206486]
- (12) Solutia study no. LF-78-144. Salmonella mutagenicity assay of O-Nitroaniline (Technical). [EPA Document no. 878211039; Fiche no. OTS0206222].
- (13) Solutia study no. ML-82-89. Orthonitroaniline: A teratology study in rats. [EPA Document no. 868600002; Fiches no. OTS0510153]
- (14) Solutia study no. ML-89-7. Micronucleus assay with o-nitroaniline.
- (15) Solutia study no. MO1983X083. Acute toxicity of o-Nitroaniline for Daphnia magna.
- (16) Solutia study no. MO20020140. Biodegradation testing of o-nitroaniline (ONA) and p-nitroaniline (PNA).
- (17) Solutia study no. Y-76-438 Toxicological investigation: O-Nitroaniline [EPA Document No. 878211634; Fiche no. OTS0206222].
- (18) Suzuki, T. 1991. J. Computer-Aided Molecular Design 5:149-166.
- (19) Zok, S, Goerge, G, Kalsch, W and Nagel, R. 1991. Sci. Total Environ. 109/110:411-421.

7.1 END POINT SUMMARY

7.2 HAZARD SUMMARY

7.3 RISK ASSESSMENT

I U C L I D

Data Set

Existing Chemical : ID: 100-01-6
CASNo. : 100-01-6
EINECS Name : 4-nitroaniline
EINECS No. : 202-810-1
TSCA Name : Benzenamine, 4-nitro-
Molecular Formula : C6H6N2O2

Producer Related Part
Company : Solutia Inc.
Creation date : 04.04.2002

Substance Related Part
Company : Solutia Inc.
Creation date : 04.04.2002

Memo :

Printing date : 07.11.2002
Revision date :
Date of last Update : 07.11.2002

Number of Pages : 43

Chapter (profile) : Chapter: 1, 2, 3, 4, 5, 7
Reliability (profile) : Reliability: without reliability, 1, 2, 3, 4
Flags (profile) : Flags: without flag, confidential, non confidential, WGK (DE), TA-Luft (DE),
Material Safety Dataset, Risk Assessment, Directive 67/548/EEC, SIDS

1.0.1 OECD AND COMPANY INFORMATION

24.10.2002

1.0.2 LOCATION OF PRODUCTION SITE

1.0.3 IDENTITY OF RECIPIENTS

1.1 GENERAL SUBSTANCE INFORMATION

1.1.0 DETAILS ON TEMPLATE

1.1.1 SPECTRA

1.2 SYNONYMS

1.3 IMPURITIES

1.4 ADDITIVES

1.5 QUANTITY

1.6.1 LABELLING

1.6.2 CLASSIFICATION

1.7 USE PATTERN

1.7.1 TECHNOLOGY PRODUCTION/USE

1.8 OCCUPATIONAL EXPOSURE LIMIT VALUES

1.9 SOURCE OF EXPOSURE

1. General Information

Id 100-01-6
Date 07.11.2002

1.10.1 RECOMMENDATIONS/PRECAUTIONARY MEASURES

1.10.2 EMERGENCY MEASURES

1.11 PACKAGING

1.12 POSSIB. OF RENDERING SUBST. HARMLESS

1.13 STATEMENTS CONCERNING WASTE

1.14.1 WATER POLLUTION

1.14.2 MAJOR ACCIDENT HAZARDS

1.14.3 AIR POLLUTION

1.15 ADDITIONAL REMARKS

1.16 LAST LITERATURE SEARCH

1.17 REVIEWS

1.18 LISTINGS E.G. CHEMICAL INVENTORIES

2.1 MELTING POINT

Value : = 146 °C
Sublimation :
Method : other
Year : 1989
GLP : no data
Test substance : other TS
Reliability : (2) valid with restrictions
Reference cited as Peer reviewed in Hazardous Substance Data Bank for p-Nitroaniline (2002) and as Recommended value in SRC CHEMFATE data base (2002).
Flag : Critical study for SIDS endpoint
07.11.2002 (1)

2.2 BOILING POINT

Value : = 332 °C at
Decomposition :
Method : other
Year : 1989
GLP : no data
Test substance : other TS
Reliability : (2) valid with restrictions
Reference cited as Peer Reviewed in Hazardous Substances Data Band for p-Nitroaniline (2002) and cited as SRC Recommended value in CHEMFATE data base (2002)
Flag : Critical study for SIDS endpoint
07.11.2002 (1)

2.3 DENSITY**2.3.1 GRANULOMETRY****2.4 VAPOUR PRESSURE**

Value : = .0053 hPa at 25° C
Decomposition :
Method : other (measured)
Year : 1985
GLP : no data
Test substance : other TS
Reliability : (2) valid with restrictions
Cited as peer reviewed reference in Hazardous Substances Data Bank for p-nitroaniline (2002).
Flag : Critical study for SIDS endpoint
24.10.2002 (3)

2.5 PARTITION COEFFICIENT

Log pow : = 1.39 at ° C

2. Physico-Chemical Data

Id 100-01-6
Date 07.11.2002

Method : other (calculated)
Year : 1987
GLP : no data
Test substance : no data
Reliability : (2) valid with restrictions
Recommended value in CHEMFATE data base (2002)
Flag : Critical study for SIDS endpoint
24.10.2002 (6)

2.6.1 WATER SOLUBILITY

Value : = 724 mg/l at 25 ° C
Qualitative :
Pka : at 25 ° C
PH : at and ° C
Method : other
Year : 1991
GLP : no data
Test substance : other TS
Reliability : (2) valid with restrictions
Cited as a Peer Reviewed reference in Hazardous Substance Data Bank
for p-nitroaniline (2002).
Flag : Critical study for SIDS endpoint
24.10.2002 (19)

2.6.2 SURFACE TENSION

2.7 FLASH POINT

2.8 AUTO FLAMMABILITY

2.9 FLAMMABILITY

2.10 EXPLOSIVE PROPERTIES

2.11 OXIDIZING PROPERTIES

2.12 ADDITIONAL REMARKS

3.1.1 PHOTODEGRADATION

Type : air
Light source : other
Light spect. : nm
Rel. intensity : based on Intensity of Sunlight
Indirect photolysis
Sensitizer : OH
Conc. of sens. :
Rate constant : = .00000000001345366 cm³/(molecule*sec)
Degradation : = 50 % after 9.5 hour(s)
Deg. Product : not measured
Method : other (calculated)
Year : 2002
GLP : no
Test substance : no data
Method : Calculated by AOP Computer Program, Vers. 1.90, Syracuse Research Corp. which estimates the Atmospheric Oxidation Potential. This program estimates the rate constant for the atmospheric, gas-phase reaction between photochemically produced hydroxyl radicals and organic chemicals. The model is based on SAR methods developed by Atkinson et al, 1987, Intern. J. Chem. Kinet. 19:799 and described in Meylan and Howard, 1993, Chemosphere 26: 2293-2299.
Reliability : (2) valid with restrictions
Estimated value based on model recommended by EPA
Flag : Critical study for SIDS endpoint
24.10.2002 (4)

3.1.2 STABILITY IN WATER

3.1.3 STABILITY IN SOIL

3.2 MONITORING DATA

3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

Type : fugacity model level III
Media : other
Air (level I) : .588
Water (level I) : 36.8
Soil (level I) : 62.6
Biota (level II / III) :
Soil (level II / III) : .0138
Method : other
Year : 2002
Method : Calculated according to Mackay, Level III. Assumed emissions (1000 kg/hr) to air, water and soil compartments using measured values as available from reference documents, including: Mol Wt=138.13; Henry's LC=1.26e-009 atm-me/mole (Henry database); Vapor Press=0.3 mm Hg (user entry); Log Kow=1.39 (user entry); Soil Koc=10.1 (calc by model). Last soil entry includes data estimate for sediments.
Results Chem Name : p-Nitroaniline
Molecular Wt: 138.13

3. Environmental Fate and Pathways

Id 100-01-6
Date 07.11.2002

Henry's LC : 1.26e-009 atm-m3/mole (Henry database)
Vapor Press : 0.3 mm Hg (user-entered)
Log Kow : 1.39 (user-entered)
Soil Koc : 10.1 (calc by model)

	Concentration (percent)	Half-Life (hr)	Emissions (kg/hr)	
Air	0.588	19	1000	
Water	36.8	20	1000	
Soil	62.6	20	1000	
Sediment	0.0138	60	0	

	Fugacity (atm)	Reaction (kg/hr)	Advection (kg/hr)	Reaction (percent)
Advection (percent)				
Air	8.89e-013	18.3	5.02	0.611
0.167				
Water	1.44e-015	1.09e+003	31.5	36.4
1.05				
Soil	5.01e-014	1.85e+003	0	61.8
0				
Sediment	2.16e-016	0.136	0.000235	0.00453
7.84e-006				

Persistence Time: 28.5 hr
Reaction Time: 28.9 hr
Advection Time: 2.34e+003 hr
Percent Reacted: 98.8
Percent Advected: 1.22

Half-Lives (hr), (based upon estimates from experimental data):

Air: 19
Water: 20
Soil: 20
Sediment: 60

Advection Times (hr):

Air: 100
Water: 1000
Sediment: 5e+004

Reliability : (2) valid with restrictions
Estimated values based on model recommended by EPA.

Flag : Critical study for SIDS endpoint

24.10.2002

(4)

3.3.2 DISTRIBUTION

3.4 MODE OF DEGRADATION IN ACTUAL USE

3.5 BIODEGRADATION

Type : aerobic
Inoculum :
Concentration : 5mg/l related to Test substance
related to
Contact time : 24 hour(s)
Degradation : = 82 % after 24 hour(s)

3. Environmental Fate and Pathways

Id 100-01-6
Date 07.11.2002

Result : other
Deg. Product :
Method : other
Year : 1975
GLP : no
Test substance : other TS
Method : Semi-continuous activated sludge (SCAS) testing was carried out over a 10-month period at an addition rate of 5 mg per 24-hr cycle. The standardized test method used was published in JAOCS 42:986 (1965) and used the modified feed technique (JAOCS 46:432, 1969). Sludge was obtained from a local waste disposal site. Disappearance was measured after one 24-hr cycle per week using UV spectrophotometry to analyze the methylene chloride extract of the mixed liquor samples taken at that time.

Result : PNA appeared to be moderately degradable under these test conditions; however, the data obtained were somewhat erratic. During the 16th through 30th week of feeding, the degradation varied from moderately rapid to rapid with a mean rate and 95% confidence limits of 82+/-12%. During the last two months of testing, far lower rates (mean of 19.4+/-10%) were observed. These data seem to indicate a threshold toxic or inhibiting effect of PNA. Substantial inhibition of the normal sludge growth rate was observed.

Test substance : Technical grade PNA with purity > 99%.
Reliability : (2) valid with restrictions
Study conducted prior to codification of GLPs but considered well documented. Methodology used has subsequently been incorporated into a standardized international test guideline for this study type.

Flag : Critical study for SIDS endpoint
07.11.2002 (16)

3.6 BOD5, COD OR BOD5/COD RATIO

3.7 BIOACCUMULATION

3.8 ADDITIONAL REMARKS

4.1 ACUTE/PROLONGED TOXICITY TO FISH

Type	:	static
Species	:	Salmo gairdneri (Fish, estuary, fresh water)
Exposure period	:	96 hour(s)
Unit	:	mg/l
Analytical monitoring	:	no
NOEC	:	= 10
LC50	:	= 45
Method	:	other
Year	:	1980
GLP	:	yes
Test substance	:	other TS
Method	:	Followed study design adopted by US EPA Committee on Methods for Toxicity Tests with Aquatic Organisms, 1975; design consistent with OECD 203. Groups of 10 fingerling (mean wt of 0.83 g/fish and length of 38 mm) were exposed to varying test concentrations in 15 liter of soft reconstituted water with a dissolved oxygen level of 8.6 mg/l, a pH of 7.4, total hardness of 45 mg/L CaCO ₃ and total alkalinity of 35 mg/l CaCO ₃ . These vessels were kept in a water bath at 12 degrees C. Fish acclimated to the dilution were held without food for 48 hours prior to testing. Based on preliminary testing, each group of fish was exposed to one of six test concentrations ranging in a logarithmic series from 5.6 to 100 mg/L. Fish were added to the test chambers within 30 min. of the addition of the test article. Test concentrations were prepared in acetone (0.5 ml), based on total compound as the test article was > 99% pure and the dose solution was then added to each respective test chamber. Mortality rates, fish behavior and water quality data (temp, pH, ammonia levels) were monitored after 24, 48 and 96 hrs of treatment. Antimycin A was similarly tested as a concurrent positive control. Calculation of the LD50 and confidence limits was performed using a computerized program developed by Stephan, Busch, Smith, Burke and Andrew, 1978 from the US EPA Duluth, Minn Aquatic Laboratory.
Result	:	LC50 and (Confidence Limits): 96-hr=46(32-56) mg/L; 48-hr= 45 (32-56) mg/L; 24-hr = 47 (32-100) mg/L. No deaths were seen at any test concentration up to 32 mg/l through 96 hrs of testing. At 56 mg/l, mortality reached 80% after 24 hrs and 90% after 48 and at 96 hrs. 100% mortality occurred at all three time points at 100 mg/l. A yellow precipitate was observed at all test levels. Dissolved oxygen concentration ranged between 60-100% saturation and was considered adequate for testing. The pH values remained consistent throughout the test and the ammonia concentrations were below the toxic limit. The positive control responded as expected.
Test substance	:	Technical grade PNA with purity > 99%.
Reliability	:	(1) valid without restriction
Flag	:	Critical study for SIDS endpoint
15.10.2002		

(9)

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Type	:	static
Species	:	Daphnia magna (Crustacea)
Exposure period	:	48 hour(s)
Unit	:	mg/l
Analytical monitoring	:	no
NOEC	:	= 10
EC50	:	= 20

Method	: other
Year	: 1980
GLP	: yes
Test substance	: other TS
Method	: Followed study design outlined by the US EPA Committee on Methods for Toxicity Tests with Aquatic Organisms, 1975, and consistent with OECD Guideline # 202. The study was conducted in 250 ml glass beakers containing 200 ml well water with specified chemical characteristics and kept at 20 degrees C. The photoperiod was controlled to give 16 hr daylight. After an initial range-find study, groups of 10 <i>D. magna</i> (first instar less than 24 hr old) were added to one of 5 beakers containing a range of test material between 3.2 and 32 mg/L, spaced logarithmically. The test article was originally prepared in 0.5 mL acetone solutions (0.5 ml) prior to charging the beakers. Each concentration was run in duplicate. Fish mortality and behavior and water quality parameters (dissolved oxygen levels, pH and temperature) were measured at the beginning of the test and after 24 hr (mortality and behavior only) and 48 hrs. Predicted LC50 values and 95% confidence limits were calculated using the computerized program developed by Stephan, Busch, Smith, Burke and Andrew, 1978 from the US EPA Duluth, Minn Aquatic Laboratory.
Result	: 48 hr LC50 (CI) =20 (18-23) mg/L. All water quality parameters (20-12 deg. C; 8.8-9.0 mg/L DO, pH of 8.1-7.9 and water hardness of 255 ppm CaCO ₃) were found to be acceptable.
Test substance	: Technical grade PNA with purity > 99%.
Reliability	: (1) valid without restriction
Flag	: Critical study for SIDS endpoint
15.10.2002	

(10)

4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

Species	: <i>Scenedesmus</i> sp. (Algae)
Endpoint	: growth rate
Exposure period	: 48 hour(s)
Unit	: mg/l
Analytical monitoring	: no data
EC50	: = 54.9
Method	: OECD Guide-line 201 "Algae, Growth Inhibition Test"
Year	: 2001
GLP	: no data
Test substance	: other TS
Method	: 48-hr algae growth inhibition test following OECD guideline 201. Organism used was <i>S. obliquus</i> . pH of the culture medium was adjusted to 7.2+/-0.2. Five concentrations were used at log intervals of 0.2. Two replicates of each concentration plus a negative control were tested. The algae in the logarithmic growing period were inoculated into 250 ml Erlenmeyer flasks containing approx 60 ml of media, test article and algae. The initial algae cell concentration was 1x10 ⁴ cells/ml. The culture was incubated under a continuous light at 20+/-1 degrees C while fluorescent lamp and the average illumination intensity was about 4000 lux. Growth was monitored by an electron microscope (400X). The EC value was determined using a one variable linear regression analysis.
Test substance	: Test material purchased from chemical supplier; typical technical grade purity of PNA was 99%.
Reliability	: (1) valid without restriction No mention was made regarding conduct under GLPs in the literature article; however, as this study was conducted specifically to meet OECD Guideline 201, it can reasonably be assumed that it also was conducted under GLPs.
Flag	: Critical study for SIDS endpoint
07.11.2002	

(7)

4.4 TOXICITY TO MICROORGANISMS E.G. BACTERIA

4.5.1 CHRONIC TOXICITY TO FISH

4.5.2 CHRONIC TOXICITY TO AQUATIC INVERTEBRATES

4.6.1 TOXICITY TO SOIL DWELLING ORGANISMS

4.6.2 TOXICITY TO TERRESTRIAL PLANTS

4.6.3 TOXICITY TO OTHER NON-MAMM. TERRESTRIAL SPECIES

4.7 BIOLOGICAL EFFECTS MONITORING

4.8 BIOTRANSFORMATION AND KINETICS

4.9 ADDITIONAL REMARKS

5.1.1 ACUTE ORAL TOXICITY

Type	: LD50
Species	: rat
Strain	: Sprague-Dawley
Sex	: male/female
Number of animals	: 25
Vehicle	: other
Value	: = 1400 mg/kg bw
Method	: other
Year	: 1976
GLP	: no
Test substance	: other TS
Method	: Consistent with # 401, but fewer animals, ie. 5 rats of mixed sex/group were given test article in 5 increasing doses at increments of 0.1 fractional log intervals; animals observed daily for 14 days for clinical signs and weighed weekly. Food and water provided ad libitum and temp./humidity controlled. Necropsies performed on all animals that died and on survivors after 14d. Technical grade PNA used, with purity > 99%. Administered as 20% solution-suspension in corn oil
Result	: OLD50 = 1400 mg/kg; Confidence Limits of 1230-1590 mg/kg; used method of deBeer, J.Pharmacol. Experimen. Ther. 85:1; Deaths - mg/kg: 794 (0/5), 1000 (1/5), 1260 (1/5), 1580 (4/5), 2000 (5/5), occurred within 7 days of dosing; Signs of toxicity: ocular discharge, tremors and convulsions; necropsy (decedents) - hemorrhagic areas of lung, liver discoloration and gi inflammation; all survivors had normal vicera after 14 days observation
Conclusion	: Sufficiently robust to provide degree of acute toxicity in rodents; numerous additional literature citations for this endpoint also available.
Reliability	: (2) valid with restrictions Conducted prior to, but consistent with, US GLPs which were enacted 6/79. Results are consistent with data in ECB IUCLID-PNA, 2002 for this endpoint, which had 5 values between 920-3250 mg/kg and 1 value as low as 750 mg/kg.
Flag	: Critical study for SIDS endpoint
07.11.2002	(17)

5.1.2 ACUTE INHALATION TOXICITY

5.1.3 ACUTE DERMAL TOXICITY

Type	: LD0
Species	: rabbit
Strain	: New Zealand white
Sex	: male/female
Number of animals	: 3
Vehicle	: other
Value	: > 7940 mg/kg bw
Method	: other
Year	: 1976
GLP	: no
Test substance	: other TS
Method	: Test article administered as 40% solution-suspension in corn oil; applied occluded for 24 hrs to intact, clipped skin of rabbits, animals observed clinically for 14 days. Body weights were recorded weekly; all animals were necropsied after d14. Food and water available ad libitum and temp./humidity was controlled.

5. Toxicity

Id 100-01-6
Date 07.11.2002

Result : temp./humidity was controlled.
: Determination of Minimum Lethal Dose: Two dosages tested, 5010 mg/kg (0/1 deaths) and 7940 mg/kg (0/2 deaths); no significant untoward toxic signs were observed during the study, all viscera normal at necropsy

Test substance Conclusion : Used Technical grade PNA, with purity of > 99%.
: Sufficiently robust study to evaluate the minimum lethal dose; as this dose proved to be of a low toxicity, there would appear to be no reason to test at higher levels to define an LD50 by this route.

Reliability : (2) valid with restrictions
This is provided as supplemental information since an acute oral toxicity study has been used to fulfill this HPV endpoint. Small, but sufficient no. animals to characterize toxicity; study conducted prior to, but consistent with, US GLPs enacted in 6/79.

07.11.2002

(17)

04.04.2002

5.1.4 ACUTE TOXICITY, OTHER ROUTES

5.2.1 SKIN IRRITATION

5.2.2 EYE IRRITATION

5.3 SENSITIZATION

5.4 REPEATED DOSE TOXICITY

Species : rat
Sex : male/female
Strain : Sprague-Dawley
Route of admin. : gavage
Exposure period : 90 days
Frequency of treatment : daily consecutive
Post obs. period : none
Doses : 0, 3, 10, 30 mg/kg/day
Control group : yes, concurrent vehicle
NOAEL : < 3 mg/kg bw
LOAEL : = 3 mg/kg bw
Method : OECD Guide-line 408 "Subchronic Oral Toxicity - Rodent: 90-day Study"
Year : 1981
GLP : yes
Test substance : other TS
Method : Corn oil vehicle used and dosing occurred at a constant volume of 0.2 ml/100 g bdy wt; 20 rats/sex/group used; Clinical signs recorded daily, individual body weights and food consumption measured weekly, serum chemistries (SGPT, SAP, BUN, T. Bili., GLU, T. Prot., K, Na), urinalysis (Prot, microscop. elements, pH, Spec. grav., blood, Glu, ketones, urobilinogen, vol.) and hematology parameters (Hgn, HCT, WBC, RBC, MCV, MCHC, retics, red cell fragility and methemoglobin) examined after 44 and 88 days. All animals necropsied at study term and organ weights (brain, adrenals, kidneys, liver, spleen, pituitary, testis) weighed. Histopathologic exams were conducted on approx. 40 tissues and organs from all high dose and control rats and the spleens of all lower dose rats. Specifically, gonads were examined for all HD and C animals. Statistical

5. Toxicity

Id 100-01-6

Date 07.11.2002

	<p>Specifically, gonads were examined for all HD and C animals. Statistical analysis performed using: Bartlett's test ($p < 0.01$), ANOVA, Dunnett's test, Mann-whitney U with Bonferroni Inequality test, and Kolmogorov-Smirnov 1 tail test (all at $p < 0.05$ and $p < 0.01$)</p>	
Result	: 30 mg/kg: Pale appearance around ears, statistically significant increase in urinary urobilinogen and methemoglobin levels, statistical increases in RBC counts and hemoglobin levels of both sexes. All animals had discolored spleens at necropsy, statistically increased spleen weights and splenomegaly and microscopic evidence of excessive splenic hemosiderin. 10 mg/kg: Statistically increased methemoglobin and decreased RBC counts and hemoglobin conc. (females only), all animals had splenomegaly, elevated splenic wts, discolored spleens and microscopic pathology associated with excessive hemosiderin; 3 mg/kg: statistically elevated methemoglobin (both sexes) and microscopic findings in spleen	
Test substance	: Used Technical grade PNA with purity > 99%.	
Conclusion	: No effects observed on gonads.	
Reliability	: (1) valid without restriction	
Flag	: Critical study for SIDS endpoint	
28.08.2002		(14)
Species	: rat	
Sex	: male/female	
Strain	: Sprague-Dawley	
Route of admin.	: inhalation	
Exposure period	: 6 hours per day, 5 days per week	
Frequency of treatment	: 4 weeks	
Post obs. period	: none	
Doses	: 0, 10, 32, 80 mg/m ³ (analytical)	
Control group	: yes, concurrent vehicle	
NOAEL	: < 10 mg/m ³	
LOAEL	: = 10 mg/m ³	
Method	: OECD Guide-line 407 "Repeated Dose Oral Toxicity - Rodent: 28-day or 14-d Study"	
Year	: 1984	
GLP	: yes	
Test substance	: other TS	
Method	: Aerosol derived by passing air over PNA dissolved in isopropanol and warmed. Groups of 10 rats/sex/group were housed in stainless steel and glass chamber and exposed under whole body conditions to one of three levels of test material. A vehicle control group was exposed to isopropanol in a similar fashion and treated similarly for evaluation. Chamber atmospheres and particle size were analytically determined. Dosing occurred 6h/d, 5d/wk for 4 consecutive weeks; animals were observed daily for clinical signs, weighed weekly, food and water given ad libitum, serum chemistry (BUN, SGPT, SAP, GLU, ALB, T.Protein, Glob., Na, K, P, Ca, Cl) and hematology (Hgb, HCT, RBC, Methem., clot time, T/Differ. Leuko, red cell morph) parameters collected on day 0 and 28. Ophthalmoscopic exams conducted on day 0 and 28. Organ weights (gonads, hrt, kid, lvr, lu, pit, spln, brain) recorded at termination; all animals necropsied at term; microscopic evaluation of approx. 40 tissues and organs (including gonads) for all high dose and control rats; spleens examined for all lower dose animals. Statistical methods used included: Bartlett's test ($p < 0.01$), and ANOVA, Kruskal-Wallis, Dunn's Summed rank test - all ($p < 0.05$ and $p < 0.01$)	
Result	: 80 mg/m ³ : non-statistical decreases in hemoglobin and hematocrit seen in males and females, statistical increase in methemoglobin in males and females, higher incidence of polychromasia and anisocytosis (females only), statistically elevated absolute and relative spleen wts for both sexes, histopathological exams revealed elevated iron deposition within splenic macrophages, extramedullary hematopoiesis in spleen (male and female) and liver (females only); 32 mg/kg: non-statistical decrease in hemoglobin in males, statistically elevated methemoglobin in males and females, higher	

5. Toxicity

Id 100-01-6
Date 07.11.2002

in males, statistically elevated methemoglobin in males and females, higher incidence of polychromasia (both sexes) and anisocytosis (females only), relative spleen wts increased statistically (males only), histopathology - increased iron deposition and extramedullary hematopoiesis in both males and females; 10 mg/m3: non-significant elevation in blood methemoglobin, significant increases in mean spleen weight (both sexes), iron deposition and extramedullary hematopoiesis seen in spleens (both sexes)

Test substance : Technical grade PNA with purity > 99%.
Reliability : (1) valid without restriction
Supplemental HPV study since a fully acceptable Subchronic study (see earlier entry in this Section) fulfills the Repeated Dose HPV Endpoint.

07.11.2002 (8)
05.04.2002

5.5 GENETIC TOXICITY 'IN VITRO'

Type : Ames test
System of testing : S. typhimurium test strains TA98, TA100, TA1535, TA1537 w & w/o S9
Concentration : 0.01, 0.04, 0.2, 1, 1.5, 3, 4, 5, and 10 mg/plate
Cycotoxic conc. : no significant microbial toxicity observed up to 10 mg/plate with TA100
Metabolic activation : with and without
Result : positive
Method : OECD Guide-line 471 "Genetic Toxicology: Salmonella typhimurium Reverse Mutation Assay"

Year : 1980
GLP : yes
Test substance : other TS
Method : Conducted both Spot test and Plate Incorporation Assay. Used DMSO as solvent, S9 was commercially available rat and mouse liver preparations. Appropriate positive (2-AA, 9-AA, B(a)P, 2-NF, NaNo2) controls run to validate method. All assays run in triplicate. Bartlett's test for homogeneity of variance and group-wise comparisons made within levels of pooled variance, 1-sided t-test applied, p<0.05. For positives, Grubb's test run to determine outliers and regression analysis and t-test of transformed data to determine dose response.

Result : Negative in all 4 test strains, with and without activation, up to max. conc. of 25 mg/spot in Spot test.
Positive finding only with TA98 (statistically elevated without activation and elevated, but not statistically with activation) in plate incorporation assay; all other strains were negative with and without activation

Test substance : Technical grade PNA with purity of > 99%.
Reliability : (1) valid without restriction
Flag : Critical study for SIDS endpoint

28.08.2002 (13)

Type : Cytogenetic assay
System of testing : Chinese Hamster Ovary cell culture
Concentration : 50 to 5000 ug/mL
Cycotoxic conc. : Laboratory 1 - 1600 ug/ml and higher; laboratory 2- none up to 5000 ug/ml
Metabolic activation : with and without
Result : ambiguous
Method : other
Year : 1986
GLP : no data
Test substance : other TS
Method : NTP study design, exposing cells for 8-12 hr normally and for 2hr in presence of S9; 100 cells per dose group were scored, all types of aberrations were recorded; Dunnett's adjusted P value (p<0.05) was used for statistical assessment.

5. Toxicity

Id 100-01-6

Date 07.11.2002

Result : Two separate testing labs used, each giving nonconfirmatory results. Positive results reported with S9 in studies at laboratory 1, and weak positive without S9 at Lab 2, Effects only seen at very highest test levels, with no evaluation of influence of pH or osmolarity. Cytotoxicity observed at Lab 1 but not reported at lab 2.

Test substance : Reportedly commercially available; i.e. technical grade of > 99%

Reliability : (3) invalid
Results considered ambiguous. Inconsistency of positive findings renders results inconclusive; additional concerns regarding inconsistency in cytotoxicity seen within lab trials and between labs. No effort made to determine affect, if any, of pH or osmolarity changes on study outcome. Supplemental HPV study since a fully acceptable in vivo micronucleus test fulfills this HPV Endpoint.

07.11.2002

(5)

Type : Cytogenetic assay

System of testing : CHO-K1 (Chinese Hamster Ovary) cells

Concentration : 173, 345, 690, and 1035 ug/ml

Cycotoxic conc. : none observed

Metabolic activation : without

Result : ambiguous

Method : other

Year : 1996

GLP : no data

Test substance : other TS

Method : Unique, research methodology performed. Used established cell line without incorporation of S9 fraction as data included in this paper considered PNA as a weak, direct acting mutagen in an Ames/Salmonella test. After incubation for 2 hrs with test compound dissolved in DMSO, cells were washed twice with PBS and incubated for another 20 hr in fresh medium. After colchicine addition, and three further hrs of incubation, metaphase cells were harvested by mitotic shake-off and resuspended. Cells were fixed, stained and selected for analysis. At least 100 metaphases per flask were scored for each dose for individual types of aberrations, including breaks, deletions, exchanges and dicentric. Both the percentage of aberrant cells and the frequency of aberrations were calculated. The tests were repeated three times in total such that at least 300 metaphases were scored for each dose. A positive response was determined based on the percentage of cells with aberrations showing a dose-response trend and at least a four-fold increase over that of the negative controls at one or more doage levels. Both Eagles' basal medium and DMSO were tested as negative controls. TEM served as a positive control.

Result : The results obtained are considered ambiguous since specified criterion for determination of a positive response (4X % aberrant cells over negative control-in this case DMSO) were not met. Neither the positive control (0.25 ug/ml TEM) nor any of the PNA dose levels exhibited a 4X increase from the negative DMSO control; the positive control and all PNA dose levels did exhibit a 4X increase in aberrant cells over the Eagle's medium negative control. The % aberrant cells reported were: Eagle's medium (3), DMSO (6), TEM (22), 173 ug/ml PNA (13), 345ug/ml PNA (19), 690 ug/ml PNA (20), and 1035 ug/ml PNA (20).

Test substance : Obtained commercially (Sigma Chem.), and thus technical grade of > 99%.

Reliability : (3) invalid
Supplemental HPV study since a fully acceptable in vivo micronucleus test is available to fulfill this endpoint; also ambiguous outcome of this study renders it unuseable.

07.11.2002

(2)

5.6 GENETIC TOXICITY 'IN VIVO'

Type	: Micronucleus assay
Species	: mouse
Sex	: male/female
Strain	: CD-1
Route of admin.	: i.p.
Exposure period	: two doses, 24-hours apart
Doses	: 80, 400 and 800 mg/kg
Result	: negative
Method	: OECD Guide-line 474 "Genetic Toxicology: Micronucleus Test"
Year	: 1987
GLP	: yes
Test substance	: other TS
Method	: High dose considered to be 80% of IP LD50, as determined by preliminary study using probit method; corn oil used as vehicle (10 ml/kg); 12 mice/sex were used for the 800 mg/kg test group, 5/sex at 400 and 80 mg/kg and 10/sex for the untreated control group; Doses were administered by IP twice with 24 hr separating each dose. Bone marrow was taken after 24 and 48 hr following last treatment from HD and C mice and after 24 h from mid and low dose animals; all mice were observed daily for clinical signs. Micronuclei recorded after assessment of 1000 PCEs/mouse at all test levels; cyclophosphamide (40 mg/kg, twice) used as positive control. Statistical significance was determined by Student's t-test (1-sided), $p < 0.05$.
Result	: No increases were seen in micronucleated PCE frequency in any PNA test group; toxicity to the cell population observed at 800 mg/kg @ 48h interval; elevated incidence of micronuclei with the positive control confirmed validity of method. One death and clear signs (unresponsiveness and tremors up to 4 hr after dosing) of toxicity were noted at 800 mg/kg; at 400 mg/kg - listlessness and some tremors seen occasionally after dosing; 80 mg/kg - listlessness immediately after dosing; No effects on body weight were observed at any test level.
Test substance	: Technical grade PNA with purity > 99%.
Reliability	: (1) valid without restriction
Flag	: Critical study for SIDS endpoint

16.10.2002

(15)

04.04.2002

5.7 CARCINOGENITY

5.8 TOXICITY TO REPRODUCTION

Type	: Two generation study
Species	: rat
Sex	: male/female
Strain	: Sprague-Dawley
Route of admin.	: gavage
Exposure period	: F0 & F1 Adults-premating through litter weaning(F0) and postweaning (F1)
Frequency of treatment	: daily (7d/wk) gavage
Premating exposure period	
Male	: F0- 14 weeks; F1 - 18 weeks
Female	: F0- 14 weeks; F1 - 18 weeks

5. Toxicity

Id 100-01-6

Date 07.11.2002

Duration of test	: F0 MF -167d; F1 MF - 216d
Doses	: 0, 0.25, 1.5 and 9 mg/kg/d
Control group	: yes, concurrent vehicle
NOAEL Parental	: >= 9 mg/kg bw
NOAEL F1 Offspr.	: >= 9 mg/kg bw
Method	: OECD Guide-line 416 "Two-generation Reproduction Toxicity Study"
Year	: 1983
GLP	: yes
Test substance	: other TS
Method	: Test material was given to groups of 15M and 30F rats (vehicle control group also included) to F0 and F1 generations during a pre-mating (14 wks for F0 and 18 wks for F1) growth period, and through the ensuing mating, gestation and lactation intervals (1 litter/generation). F1 rats continued on treatment during a post-weaning period of 30d. Dosing concentrations confirmed for accuracy. Body weights were recorded weekly for F0 and F1M. For F0 and F1 F wts were recorded weekly through the growth period and up to mating, then resumed after mating until sacrifice. Food consumption was recorded weekly for F0 and F1 M from study start up to mating, then resumed after mating through study term. Food consumption for adult females F0 and F1 was recorded weekly through the growth period and again after weaning of litters. Cageside observations were made weekly, as well as daily observations of clinical signs. Temperature, humidity and light-dark cycles were controlled. F0 adults were sacrificed following weaning of the F1 litters and given a gross postmortem examination; reproductive tissues (testes, epididymides, seminal vesicles) were evaluated histopathologically for all control and high dose males. Adult M and F rats were sacrificed following completion of a post-weaning treatment interval, given a gross necropsy, and full histopathological examination of over 40 tissues and organs (including gonads) performed on 10 randomly selected animals/sex/group. Pups delivered to F0 and F1 females were evaluated for growth, survival and external irregularities during lactation days 0, 4, 14 and 21. F1 pups not selected for the adult generation were sacrificed and given a gross postmortem exam. Tissues were evaluated histopathologically (~40 tissues/organs) from 5/sex/group of F1 pups.
Result	: No adverse effects observed in either F0 or F1 adults in mortality, body weights or food consumption or physical in-life evaluations. Mating indices were comparable to controls for both F0 and F1. A statistically significant reduction in pregnancy rate was observed in the 9 mg/kg F0 group vs concurrent control value, and just outside of laboratory historical control range. The male fertility index was slightly, but not statistically, lower at 9 mg/kg dose in F0. Both male and female fertility indices in F1 generation were comparable to control group at all test levels. No adverse effects were observed in mean length of gestation, no. live and dead pups at monitored time points, pup weights during lactation, pup and litter survival. No compound-related gross postmortem changes were observed during examination of any F0 or F1 adults or offspring. No microscopic changes were noted with respect to gonads evaluated on F0 adults or F1 offspring.
Test substance	: Technical grade PNA with purity > 99%.
Conclusion	: The reduction in female fertility index seen in F0 adults is considered unrelated to treatment for the following reasons: No similar findings occurred in F1 Females, even though they were exposed for a substantially longer period (both in utero and during pre-mating phase) than their F0 counterparts and there was no evidence of histological changes in gonads which could account for this finding; Similarly, no treatment-related effects were observed on the gonads of rats exposed for up to 2 years by the same dosage (9 mg/kg/d) by the same exposure route (gavage) (Nair et al FAAT 15:607-621)
Reliability	: (1) valid without restriction
Flag	: Critical study for SIDS endpoint

16.10.2002

(11)

5.9 DEVELOPMENTAL TOXICITY/TERATOGENICITY

Species	:	rat
Sex	:	female
Strain	:	Sprague-Dawley
Route of admin.	:	gavage
Exposure period	:	gestation days 6 through 19
Frequency of treatment	:	once per day, gestation days 6-19
Duration of test	:	dosing during gestation days 6-19, sacrificed on day 20
Doses	:	0, 25, 85, 250 mg/kg
Control group	:	yes, concurrent vehicle
NOAEL Maternalt.	:	= 25 mg/kg bw
NOAEL Teratogen	:	= 85 mg/kg bw
NOAEL Embryotoxicity	:	= 85 mg/kg bw
NOAEL Fetotoxicity	:	= 25 mg/kg bw
Method	:	OECD Guide-line 414 "Teratogenicity"
Year	:	1980
GLP	:	yes
Test substance	:	other TS
Method	:	24 pregnant female rats per group; dosing occurred during days 6-19; vehicle used was corn oil (10 ml/kg constant volume), Corn oil vehicle control also included. Nidation data collected at sacrifice; live fetuses examined externally and by Wilson sections and skeletal exam techniques used to detect any variations or abnormalities. Body weights collected on gestation days 3, 6, 8, 13, 15, 17 and 20. Statistical methods used: body wts analyzed using Dunnett's test, Counted data (corpora lutea, implants, resorptions, live/dead pups) were analyzed using Mann-whitney U test; Response data (eg. pregnancy rates, litters with postimplantation loss, etc.) assessed with Fischer's exact test and Chi square test. (p<0.05 and p<0.01),
Remark	:	Supplemental information for HPV program as an adequate 2-generation study is available on PNA to fulfill the Reproductive Toxicity Endpoint.
Result	:	250 mg/kg: Reduced maternal wt gain between d6-d20, observations - pale eye coloration and occasional convulsions after dosing, significant increase in mean no. resorptions and % resorptions, significant increase in maternal mean spleen wts (abs. and rel), significantly lower mean fetal wts (both sexes), significant increase in no. fetuses with ossif. variations and fetuses with external, soft tissue or skeletal malformations (predominantly kinked or shortened tail, absence of kidneys or ureter and fused ribs); 85 mg/kg - Significant increase in mean maternal spleen wts, significantly lower mean fetal wts (both sexes); no increases in variations or malformations; 25 mg/kg - no effects on maternal, embryo- or fetotoxicity and no increase in malformations; 25 mg/kg - no treatment-related effects on maternal, embryotoxicity, fetotoxicity or terata.
Test substance	:	Technical grade PNA with purity > 99%.
Reliability	:	(1) valid without restriction
28.08.2002		

(18)

Species	:	rabbit
Sex	:	female
Strain	:	New Zealand white
Route of admin.	:	gavage
Exposure period	:	gestation days 7 through 19
Frequency of treatment	:	daily
Duration of test	:	dosed from gestation day 7 through 19, sacrificed on g. day 30
Doses	:	0, 15, 75, 125 mg/kg
Control group	:	yes, concurrent vehicle
NOAEL Maternalt.	:	= 75 mg/kg bw
NOAEL Teratogen	:	= 125 mg/kg bw

5. Toxicity

Id 100-01-6

Date 07.11.2002

NOAEL Embryotoxicity	:	= 125 mg/kg bw
NOAEL Fetotoxicity	:	= 125 - mg/kg bw
Method	:	OECD Guide-line 414 "Teratogenicity"
Year	:	1981
GLP	:	yes
Test substance	:	other TS
Method	:	18 mated females used per dose group; vehicle used was corn oil. Treated and control groups (corn oil) were dosed at constant volume of 2 ml/kg; Observations made for signs of toxicity on gestation days 0, 7, 10, 15, 19, 25 and 30; Body weights recorded on gestation days 0, 7, 19 and 30. Nidation data collected at sacrifice (gestation day 30). live fetuses examined externally and by Wilson sections and skeletal exam techniques to detect any variations or abnormalities. Statistical methods used: Bartlett's and ANOVA, Dunnett's test, Mann-whitney U test, Dunn's Rank Sum, Fischer's exact test and Jonckheere's test; $p < 0.05$ and $p < 0.01$.
Remark	:	Supplemental information for HPV program as an adequate 2-generation study is available on PNA to fulfill the Reproductive Toxicity Endpoint.
Result	:	125 mg/kg - 7/18 deaths between gestation days 14 and 20, observations - grayish appearing eyes; overall body wt gain similar to controls but higher no. of animals which lost wt during dosing observed at this test level; no increase in absol or rel spleen wt; incidence of spontaneous abortions was 4 (vs 2 for controls), however, this incidence level was frequently seen with rabbits at the test facility and thus could not be attributed to test article; no significant differences observed in mean no. implantations, resorptions or viable fetues or mean fetal wts between treated and control group; incidence and types of ossification variations in fetuses, soft tissue anomalies and external malformations were similar between treated and control groups; a slightly higher (not statistically significant) incidence in skeletal malformations was observed in treated groups vs. controls but was not considered treatment related as there was no dose response relationship for individual malformations identified in this study and they have been observed as spontaneous lesions in this rabbit strain; 75 mg/kg: observations - grayish eyes, otherwise no effects on other measured maternal, embryo, or fetal parameters. No evidence of treatment-related effect on variations or malformations; 15 mg/kg - no treatment related study findings
Test substance	:	Technical grade PNA with purity of > 99%.
Reliability	:	(1) valid without restriction

07.11.2002

(12)

05.04.2002

05.04.2002

5.10 OTHER RELEVANT INFORMATION

5.11 EXPERIENCE WITH HUMAN EXPOSURE

- (1) Budavari, S. (ed.) 1989. The Merck Index. - an encyclopedia of chemicals, drugs and biologicals, Whitehouse, NY. p.1042.
- (2) Chung, K-T, Murdock, CA, Yanguai, Z, Stevens, Jr, SE, Li, Y-S, Wei, C-I, Fernando, SY and Chou, M-W. 1996. Effects of the nitro-group on the mutagenicity and toxicity of some benzamines. Environ Molec. Mutagen. 27:67-74.
- (3) Dixon and Rissmann. 1985.
- (4) EPIWIN, version 3.10. 2002. Syracuse Research Corp., Syracuse, NY.
- (5) Galloway, SM, Armstrong, MJ, Reuben, C, Colman, S, Brown, B, Cannon, C, Bloom, AD, Nakamura, F, Ahmed, M, Duk, S, Rimpo, J, Margolin, BH, Resnick, MA, Anderson, B, and Zeiger, E. 1987. Environ Molec Mutagenesis 10:1-175..
- (6) Hansch and Leo, 1987. The Log P Database. Claremont, Ca: Pomona College.
- (7) Lu, G.-H, Yuan, X, and Zhao, Y-H. 2001. Chemosphere 44:437-440.
- (8) Solutia study BD-80-508. A four-week inhalation toxicity study of para-Nitroaniline in the rat; also published as Nair et al. (1986) Fundamen Appl Toxicol 6:618
- (9) Solutia study no. AB-80-318. Acute toxicity of para-Nitroaniline to Rainbow Trout.
- (10) Solutia study no. AB-80-320. Acute toxicity of p-Nitroaniline to Daphnia magna.
- (11) Solutia study no. BD-80-471. A 2-generation reproduction study of p-Nitroaniline in rats; also published as Nair, RS et al. 1990. FAAT 6:607-621.
- (12) Solutia study no. BD-80-529; also published as Nair, RS et al 1985 in Toxicology of Nitroaromatic Compounds. Hemisphere Publishing, NY.
- (13) Solutia study no. DA-79-257. Salmonella mutagenicity assay of CP 26926 (p-Nitroaniline). [EPA Documentation No. 878211039; Fiche No. OTS0206222].
- (14) Solutia Study no. ML-79-11. Ninety-day study of p-Nitroaniline administered to male and female Sprague-Dawley rats via gavage; also Houser, TAP 3:128
- (15) Solutia study no. ML-87-8. Mouse micronucleus assay with p-Nitroaniline.
- (16) Solutia study no. MO20020140. Biodegradation testing of o-Nitroaniline and p-Nitroaniline.
- (17) Solutia study no. Y-76-35. Toxicological Investigaion: P-Nitroaniline.
- (18) Study no. BD-79-326; also published as Nair, RS et al 1985 in Toxicology of Nitroaromatic Compounds. Hemisphere Publishing, NY.
- (19) Suzuki. J. 1991. J. Computer-Aided Molecular Design 5: 149-166.

7.1 END POINT SUMMARY

7.2 HAZARD SUMMARY

7.3 RISK ASSESSMENT